



Master thesis: Polarization-controlled active fiber-based retroreflector for laser spectroscopy of atomic deuterium

Laser spectroscopy division (Prof. T. W. Hänsch), Max Planck Institute of Quantum Optics

Precision laser spectroscopy of atomic hydrogen and deuterium has long been used to determine physical constants and to test fundamental theories, owing to the simple structure of atoms allowing comparison with precise calculations. We are continuously improving constraints on these tests and shine light on possible discrepancies, such as the proton radius puzzle [1]. At the moment, we are measuring the 2S-6P transition in deuterium which allows for determination of the deuteron radius and further tests of quantum electrodynamics. To reach the required fractional uncertainty of a few parts in 10^{12} of the transition frequency, a sophisticated control over systematic effects such as quantum interference is required [2].

Though similar to a hydrogen measurement, the spectroscopy of deuterium requires careful control of the laser polarization and a detailed understanding of the atomic line shape. We are **looking for a master student for the deuterium spectroscopy experiment** who is interested in working with a typical atomic physics toolkit, including **optics, lasers, fibers and electronics**. More specifically, the master student would investigate imperfect polarization in our improved active fiber-based retroreflector [3] and explore possibilities to extend the setup for circular polarization. Furthermore, a fiber-based polarization-controller could be used to stabilize polarization. Another aspect is the detailed investigation of polarization properties of single-mode optical fibers, along with a thorough study of fiber parameters. Though theoretical understanding and simulations of how imperfect polarization enters the deuterium spectroscopy could be also part of the thesis, the project would include mostly hands-on experimental work.

If you are interested in joining our team and want to learn more about the project, do not hesitate to contact Vitaly Wirthl (vitaly.wirthl@mpq.mpg.de). Please send your application along with your CV to Prof. Thomas Udem (thu@mpq.mpg.de) with CC to Vitaly.




Photo showing the vacuum chamber used to measure the 2S-4P transition frequency in atomic hydrogen. The purple glow in the back stems from the microwave discharge that dissociates hydrogen molecules into hydrogen atoms. The blue light in the front is fluorescence of the vacuum viewport from the ultraviolet laser that excites the atoms to the 2S state. The turquoise blue glow is stray light from the laser system used to measure the frequency of the 2S-4P transition. (Credit: Axel Beyer / MPQ)

Key references

- [1] A. Beyer *et al.* The Rydberg constant and proton size from atomic hydrogen. *Science* **358**, 79–85 (2017).
- [2] Th. Udem *et al.* Quantum Interference Line Shifts of Broad Dipole-Allowed Transitions. *Annalen der Physik* **531**, 1900044 (2019).
- [3] V. Wirthl *et al.* Improved active fiber-based retroreflector with intensity stabilization and a polarization monitor for the near UV. *Opt. Express* **29**(5), 7024-7048 (2021).