

# Pair production in counter-propagating laser pulses

JG Kirk<sup>1</sup> AR Bell<sup>2,3</sup> & I Arka<sup>1</sup>

<sup>1</sup> Max-Planck-Institut für Kernphysik, Postfach 10 39 80, 69029 Heidelberg, Germany

<sup>2</sup> Clarendon Laboratory, University of Oxford, Parks Road, Oxford OX1 3PU, UK

<sup>3</sup> Central Laser Facility, Rutherford Appleton Laboratory, Oxfordshire, OX11 0QX, UK

*E-mail: john.kirk@mpi-hd.mpg.de*

**Abstract:** We study the classical trajectories of electrons in an under-dense plasma in the presence of intense, counter-propagating laser pulses. By computing the photon emissivity and the photon trajectories through a cylindrical model of the intersecting beams we evaluate the local production rate of electron positron pairs via the non-linear Breit-Wheeler process.

Pair production has been reported in several experiments that use intense laser beams either as an accelerator or as a target for electron beams. In counter-propagating beams, however, the laser can simultaneously act as both an accelerator and a target [1]. In beams of intensity  $10^{23}$ – $10^{24}$  Wcm<sup>-2</sup>, electrons radiate gamma rays, which can produce electron-positron pairs on interacting with the laser fields via the non-linear Breit-Wheeler process. The pairs, in their turn, will be accelerated, leading to an electromagnetic cascade that could deplete much of the beam's energy [2]. We present the gamma-ray spectrum and the expected pair yield per electron in realistically modeled laser pulse-trains of both circular and linear polarisation.

## References

- [1] A.R. Bell, J.G. Kirk “Possibility of prolific pair production with high-power lasers” Phys. Rev. Lett. *101*, 200403 (2008).
- [2] J.G. Kirk, A.R. Bell, I. Arka “Pair production in counter-propagating laser beams” Plasma Physics and Controlled Fusion *51*, 085008 (2009).